

## WHAT TO EXPECT POST YOUR UROLIFT PROCEDURE

This information explains what you may experience post your procedure and how to take care of yourself during this time.

### **Expect:**

- Patients may experience some urinary discomfort during the recovery period
- Blood in the urine which can last up to 2-4 weeks post procedure
- Pain or discomfort when urinating, increased urge and frequency in urination which can last up to 4 weeks
- The catheter is generally removed on the morning after surgery. The expected hospital stay is no more than one night
- If you have had an anaesthetic you will be uninsured to drive for 24hrs
- Return to work is usually 1 week post procedure but this depends on how your recovery is going, type of work and when you are comfortable. Dr Elmes rooms can provide you with a medical certificate if required.

#### What to do:

- · Drink plenty of fluids
- Take **Ural** for the first 2-3 days if you have penile pain during urination. This helps make your urine more like water and less like lemon juice (maximum sachets a day is 4)
- Movicol is available from your local pharmacy used to avoid constipation
- Oxytrol OR Vesicare \*DO NOT take these two medications at the same time

# OXYTROL – is a skin patch for the treatment of overactive bladder with symtoms of urge urinary incontinence, urgency, and frequency. It delivers the active ingredient directly into your blood stream through your skin.

- Stick a patch anywhere on your skin, replace every 3-4 days. This is used for if you have bothersome urgency and frequency, night time urination incontinence/or penile tip pain
- These can be used a maximum of 2 patches at any one time
- Side effects: dry mouth, constipation, dry eyes, forgetfulness and rarely urinary retention

# <u>VESICARE</u> - is a tablet form to treat overactive bladder with symptoms of urgency, <u>frequency</u>, and <u>leakage</u>

- Side effects: dry mouth, constipation, dry eyes, forgetfulness and rarely urinary retention
- **Endone** prescribed strong pain relieving drug. *Side effects:* dry mouth, constipation, nausea/vomiting, dizziness, light headed (do not drive if affected)

### Avoid:

- Straining/strenuous activity and constipation for 4 weeks
- Driving long distance within first week

### Inform Dr Elmes' rooms if:

- Unable to urinate
- Heavy continuous bleeding / multiple clots
- Fevers / unwell