



Dr Martin Elmes
Urological Surgeon
Gold Coast Urologist
P: 07 5575 7922
Suite 2 Level 1, Varsity Lakes Day Hospital,
2 Lake Street Varsity Lakes 4227

WHAT TO EXPECT POST YOUR HOLEP PROCEDURE

This information explains what you may experience post your procedure and how to take care of yourself during this time.

Expect:

- There is always some bleeding from the prostate after the procedure. Blood in the urine can last up to 2-4 weeks post your procedure
- The catheter is generally removed on the morning after surgery. The expected hospital stay is no more than one night
- Return to work can take up to 2 weeks post procedure but this depends on how your recovery is going, type of work and when you are comfortable. Dr Elmes rooms can provide you with a medical certificate if required.

What to do:

- **Drink plenty of fluid**
- Take **Ural** for the first 2-3 days if you have penile pain during urination. This helps make your urine more like water and less like lemon juice (maximum sachets a day is 4)
- **Movicol** is available from your local pharmacy – used to avoid constipation
- **Oxytrol OR Vesicare**

***DO NOT take these two medications at the same time**

OXYTROL – is a skin patch for the treatment of overactive bladder with symptoms of urge urinary incontinence, urgency, and frequency. It delivers the active ingredient directly into your blood stream through your skin.

- Stick a patch anywhere on your skin, replace every 3-4 days. This is used for if you have bothersome urgency and frequency, night time urination incontinence/or penile tip pain
- These can be used a maximum of 2 patches at any one time
- Side effects: dry mouth, constipation, dry eyes, forgetfulness and rarely urinary retention

VESICARE - is a tablet form to treat overactive bladder with symptoms of urgency, frequency, and leakage

- Side effects: dry mouth, constipation, dry eyes, forgetfulness and rarely urinary retention
- **Endone** – prescribed strong pain relieving drug. *Side effects:* dry mouth, constipation, nausea/vomiting, dizziness, light headed (do not drive if affected)

Avoid:

- Straining/strenuous activity and constipation for 4 weeks
- Driving long distance within first week

Inform Dr Elmes' rooms if:

- Unable to urinate
- Heavy continuous bleeding +/- clots
- Fevers/unwell