

Dr Martin Elmes  
Urological Surgeon - Gold Coast Urologist  
P: 07 5575 7922  
Suite 2 Level 1, Varsity Lakes Day Hospital,  
2 Lake Street Varsity Lakes 4227



UROLOGICAL SURGEON  
DR MARTIN ELMES  
MBBS (MELB), FRACS (UROL. SYD)

## WHAT TO EXPECT POST YOUR HOLEP PROCEDURE

This information explains what you may experience post your procedure and how to take care of yourself during this time.

### **Expect:**

- There is usually some bleeding from the prostate after the procedure. Blood in the urine can last up to 6 weeks post your procedure, but usually settles at 2 weeks
- The catheter is generally removed on the morning after surgery. The expected hospital stay is usually one night
- Return to work can take up to 2 weeks post procedure but this depends on how your recovery is going, type of work and when you are comfortable. Dr Elmes rooms can provide you with a medical certificate if required.

### **What to do:**

- Drink plenty of fluid
- Take **Ural** for the first 2-3 days if you have penile pain during urination. This helps make your urine more like water and less like lemon juice (maximum sachets a day is 4)
- **Movicol** is available from your local pharmacy – used to avoid constipation

### **Frequency/urgency/penile tip pain**

- **Oxytrol patches OR Vesicare** is used to assist. These can be used a maximum of 2 patches/or tablets at any one time. Side effects: dry mouth, constipation, dry eyes, forgetfulness and rarely urinary retention. DO NOT use these two medications at the same time unless instructed by Dr Elmes.

### **Incontinence**

- 20% of men experience leakage to varying degrees. You should utilise either Oxytrol patches or Vesicare (never both at the same time unless instructed by Dr Elmes) as well as avoid caffeine/alcohol
- Pelvic floor exercises are recommended to build your pelvic floor muscles and control your bladder post operatively. Your pelvic floor muscle can be found during urination when you cut off your flow. The muscle you use to do this is what you are training and strengthening. Try to utilise this muscle whenever you think of it. You can also use incontinence pads from your local pharmacy if necessary.

### **Pain**

- Panadine Forte/Endone/Targin/Tapentadol – prescribed strong pain relieving drug. Side effects: dry mouth, constipation, nausea/vomiting, dizziness, light headed (do not drive if affected)

### **Avoid:**

- Straining/strenuous activity and constipation for 4 weeks (this includes the gym)

- Driving within first week, thereafter drive cautiously
- Sexual activity 6 weeks post operatively

**Inform Dr Elmes' rooms if:**

- Unable to urinate
- Heavy continuous bleeding +/- clots
- Fevers/unwell